


# Atoms, Molecules and Ions

## Chapter 2




Copyright © The McGraw-Hill Companies, Inc. Permission required for reproduction or display.

## Dalton's Atomic Theory (1808)


1. Elements are composed of extremely small particles called **atoms**.
2. All **atoms** of a given element are identical, having the same size, mass and chemical properties. The atoms of one element are different from the atoms of all other elements.
3. **Compounds** are composed of atoms of more than one element. In any compound, the ratio of the numbers of atoms of any two of the elements present is either an integer or a simple fraction.
4. A **chemical reaction** involves only the separation, combination, or rearrangement of atoms; it does not result in their creation or destruction.

## Dalton's Atomic Theory

Carbon monoxide




Carbon dioxide




Law of Multiple Proportions


3



Atoms of element X



Atoms of element Y





Compounds of elements X and Y




Law of Conservation of Mass

4




### Millikan's Experiment



**Measured mass of  $e^-$**   
(1923 Nobel Prize in Physics)

$e^-$  charge =  $-1.60 \times 10^{-19}$  C  
Thomson's charge/mass of  $e^-$  =  $-1.76 \times 10^8$  C/g  
 $e^-$  mass =  $9.10 \times 10^{-28}$  g 7

### Types of Radioactivity




Lead block  
Radioactive substance  
(uranium compound)

8

## Thomson's Model


Positive charge spread  
over the entire sphere



9

## Rutherford's Experiment


(1908 Nobel Prize in Chemistry)




1. atoms positive charge is concentrated in the nucleus
2. proton (p) has opposite (+) charge of electron (-)
3. mass of p is 1840 x mass of  $e^-$  ( $1.67 \times 10^{-24}$  g)

10

### Rutherford's Model of the Atom




atomic radius  $\sim 100 \text{ pm} = 1 \times 10^{-10} \text{ m}$   
 nuclear radius  $\sim 5 \times 10^{-3} \text{ pm} = 5 \times 10^{-15} \text{ m}$



“If the atom is the Houston Astrodome, then the nucleus is a marble on the 50-yard line.” 11

### Chadwick's Experiment (1932) (1935 Noble Prize in Physics)

H atoms - 1 p; He atoms - 2 p  
 mass He/mass H should = 2  
 measured mass He/mass H = 4



neutron (n) is neutral (charge = 0)

n mass  $\sim$  p mass =  $1.67 \times 10^{-24} \text{ g}$

12

**TABLE 2.1** Mass and Charge of Subatomic Particles

Particle	Mass (g)	Charge	
		Coulomb	Charge Unit
Electron*	$9.10938 \times 10^{-28}$	$-1.6022 \times 10^{-19}$	-1
Proton	$1.67262 \times 10^{-24}$	$+1.6022 \times 10^{-19}$	+1
Neutron	$1.67493 \times 10^{-24}$	0	0

\*More refined measurements have given us a more accurate value of an electron's mass than Millikan's.

mass p  $\approx$  mass n  $\approx$  1840 x mass e<sup>-</sup>


13

## Atomic number, Mass number and Isotopes

**Atomic number** (Z) = number of protons in nucleus


**Mass number** (A) = number of protons + number of neutrons  
= atomic number (Z) + number of neutrons

**Isotopes** are atoms of the same element (X) with different numbers of neutrons in their nuclei



14

## The Isotopes of Hydrogen



15

How many protons, neutrons, and electrons are in  ${}^{14}_6\text{C}$ ?

6 protons, 8 (14 – 6) neutrons, 6 electrons

How many protons, neutrons, and electrons are in  ${}^{11}_6\text{C}$ ?

6 protons, 5 (11 – 6) neutrons, 6 electrons

16



### The Modern Periodic Table

Period

Group

Halogen

Noble Gas

Alkali Metal

Alkali Earth Metal

Metals


Metalloids

Nonmetals

58	59	60	61	62	63	64	65	66	67	68	69	70	71
Ce	Pr	Nd	Pm	Sm	Eu	Gd	Tb	Dy	Ho	Er	Tm	Yb	Lu
90	91	92	93	94	95	96	97	98	99	100	101	102	103
Th	Pa	U	Np	Pu	Am	Cm	Bk	Cf	Es	Fm	Md	No	Lr

17

A **molecule** is an aggregate of two or more atoms in a definite arrangement held together by chemical forces



A **diatomic molecule** contains only two atoms

$\text{H}_2$ ,  $\text{N}_2$ ,  $\text{O}_2$ ,  $\text{Br}_2$ ,  $\text{HCl}$ ,  $\text{CO}$

diatomic elements


A **polyatomic molecule** contains more than two atoms

$\text{O}_3$ ,  $\text{H}_2\text{O}$ ,  $\text{NH}_3$ ,  $\text{CH}_4$

An **ion** is an atom, or group of atoms, that has a net positive or negative charge.



**cation** – ion with a positive charge

If a neutral atom **loses** one or more electrons it becomes a cation.




**anion** – ion with a negative charge


If a neutral atom **gains** one or more electrons it becomes an anion.



A **monatomic ion** contains only one atom



A **polyatomic ion** contains more than one atom



## Common Ions Shown on the Periodic Table

1 1A	2 2A	3 3B	4 4B	5 5B	6 6B	7 7B	8 8B	9 8B	10 8B	11 1B	12 2B	13 3A	14 4A	15 5A	16 6A	17 7A	18 8A
Li <sup>+</sup>													C <sup>4-</sup>	N <sup>3-</sup>	O <sup>2-</sup>	F <sup>-</sup>	
Na <sup>+</sup>	Mg <sup>2+</sup>											Al <sup>3+</sup>		P <sup>3-</sup>	S <sup>2-</sup>	Cl <sup>-</sup>	
K <sup>+</sup>	Ca <sup>2+</sup>				Cr <sup>2+</sup> Cr <sup>3+</sup>	Mn <sup>2+</sup> Mn <sup>3+</sup>	Fe <sup>2+</sup> Fe <sup>3+</sup>	Co <sup>2+</sup> Co <sup>3+</sup>	Ni <sup>2+</sup> Ni <sup>3+</sup>	Cu <sup>+</sup> Cu <sup>2+</sup>	Zn <sup>2+</sup>				Se <sup>2-</sup>	Br <sup>-</sup>	
Rb <sup>+</sup>	Sr <sup>2+</sup>									Ag <sup>+</sup>	Cd <sup>2+</sup>		Sn <sup>2+</sup> Sn <sup>4+</sup>		Te <sup>2-</sup>	I <sup>-</sup>	
Cs <sup>+</sup>	Ba <sup>2+</sup>									Au <sup>+</sup> Au <sup>3+</sup>	Hg <sub>2</sub> <sup>2+</sup> Hg <sup>2+</sup>		Pb <sup>2+</sup> Pb <sup>4+</sup>				

21









How many protons and electrons are in  ${}_{13}^{27}\text{Al}^{3+}$  ?

13 protons, 10 (13 – 3) electrons

How many protons and electrons are in  ${}_{34}^{78}\text{Se}^{2-}$  ?

34 protons, 36 (34 + 2) electrons



22

Formulas and Models				
	Hydrogen	Water	Ammonia	Methane
Molecular formula	$H_2$	$H_2O$	$NH_3$	$CH_4$
Structural formula	$H-H$	$H-O-H$	$\begin{array}{c} H-N-H \\   \\ H \end{array}$	$\begin{array}{c} H \\   \\ H-C-H \\   \\ H \end{array}$
Ball-and-stick model				
Space-filling model				

23

A **molecular formula** shows the exact number of atoms of each element in the smallest unit of a substance




An **empirical formula** shows the simplest whole-number ratio of the atoms in a substance

**molecular****empirical**

24



## Formula of Ionic Compounds



27

## Chemical Nomenclature

- **Ionic Compounds**

- Often a metal + nonmetal
- Anion (nonmetal), add “ide” to element name



barium chloride



potassium oxide



magnesium hydroxide



potassium nitrate

28

- Transition metal ionic compounds
  - indicate charge on metal with Roman numerals

A simplified periodic table with the transition metal blocks highlighted in green. The blocks are labeled 3B, 4B, 5B, 6B, 7B, 8B, and 1B, 2B.

$\text{FeCl}_2$     2  $\text{Cl}^-$   $-2$  so Fe is  $+2$         iron(II) chloride

$\text{FeCl}_3$     3  $\text{Cl}^-$   $-3$  so Fe is  $+3$         iron(III) chloride

$\text{Cr}_2\text{S}_3$     3  $\text{S}^{2-}$   $-6$  so Cr is  $+3$  ( $6/2$ ) chromium(III) sulfide

29

**TABLE 2.2** The “-ide” Nomenclature of Some Common Monatomic Anions According to Their Positions in the Periodic Table

Group 4A	Group 5A	Group 6A	Group 7A
C carbide ( $\text{C}^{4-}$ )*	N nitride ( $\text{N}^{3-}$ )	O oxide ( $\text{O}^{2-}$ )	F fluoride ( $\text{F}^-$ )
Si silicide ( $\text{Si}^{4-}$ )	P phosphide ( $\text{P}^{3-}$ )	S sulfide ( $\text{S}^{2-}$ )	Cl chloride ( $\text{Cl}^-$ )
		Se selenide ( $\text{Se}^{2-}$ )	Br bromide ( $\text{Br}^-$ )
		Te telluride ( $\text{Te}^{2-}$ )	I iodide ( $\text{I}^-$ )

\*The word “carbide” is also used for the anion  $\text{C}_2^{2-}$ .

30

**TABLE 2.3** Names and Formulas of Some Common Inorganic Cations and Anions

Cation	Anion
aluminum ( $\text{Al}^{3+}$ )	bromide ( $\text{Br}^-$ )
ammonium ( $\text{NH}_4^+$ )	carbonate ( $\text{CO}_3^{2-}$ )
barium ( $\text{Ba}^{2+}$ )	chlorate ( $\text{ClO}_3^-$ )
cadmium ( $\text{Cd}^{2+}$ )	chloride ( $\text{Cl}^-$ )
calcium ( $\text{Ca}^{2+}$ )	chromate ( $\text{CrO}_4^{2-}$ )
cesium ( $\text{Cs}^+$ )	cyanide ( $\text{CN}^-$ )
chromium(III) or chromic ( $\text{Cr}^{3+}$ )	dichromate ( $\text{Cr}_2\text{O}_7^{2-}$ )
cobalt(II) or cobaltous ( $\text{Co}^{2+}$ )	dihydrogen phosphate ( $\text{H}_2\text{PO}_4^-$ )
copper(I) or cuprous ( $\text{Cu}^+$ )	fluoride ( $\text{F}^-$ )
copper(II) or cupric ( $\text{Cu}^{2+}$ )	hydride ( $\text{H}^-$ )
hydrogen ( $\text{H}^+$ )	hydrogen carbonate or bicarbonate ( $\text{HCO}_3^-$ )
iron(II) or ferrous ( $\text{Fe}^{2+}$ )	hydrogen phosphate ( $\text{HPO}_4^{2-}$ )
iron(III) or ferric ( $\text{Fe}^{3+}$ )	hydrogen sulfate or bisulfate ( $\text{HSO}_4^-$ )
lead(II) or plumbous ( $\text{Pb}^{2+}$ )	hydroxide ( $\text{OH}^-$ )
lithium ( $\text{Li}^+$ )	iodide ( $\text{I}^-$ )
magnesium ( $\text{Mg}^{2+}$ )	nitrate ( $\text{NO}_3^-$ )
manganese(II) or manganous ( $\text{Mn}^{2+}$ )	nitride ( $\text{N}^{3-}$ )
mercury(I) or mercurous ( $\text{Hg}_2^{2+}$ )*	nitrite ( $\text{NO}_2^-$ )
mercury(II) or mercuric ( $\text{Hg}^{2+}$ )	oxide ( $\text{O}^{2-}$ )
potassium ( $\text{K}^+$ )	permanganate ( $\text{MnO}_4^-$ )
rubidium ( $\text{Rb}^+$ )	peroxide ( $\text{O}_2^{2-}$ )
silver ( $\text{Ag}^+$ )	phosphate ( $\text{PO}_4^{3-}$ )
sodium ( $\text{Na}^+$ )	sulfate ( $\text{SO}_4^{2-}$ )
strontium ( $\text{Sr}^{2+}$ )	sulfide ( $\text{S}^{2-}$ )
tin(II) or stannous ( $\text{Sn}^{2+}$ )	sulfite ( $\text{SO}_3^{2-}$ )
zinc ( $\text{Zn}^{2+}$ )	thiocyanate ( $\text{SCN}^-$ )

\*Mercury(I) exists as a pair as shown.

31

## • Molecular compounds

- Nonmetals or nonmetals + metalloids
- Common names
  - $\text{H}_2\text{O}$ ,  $\text{NH}_3$ ,  $\text{CH}_4$ ,
- Element furthest to the left in a period and closest to the bottom of a group on periodic table is placed first in formula
- If more than one compound can be formed from the same elements, use prefixes to indicate number of each kind of atom
- Last element name ends in *ide*

**TABLE 2.4**

Greek Prefixes Used in Naming Molecular Compounds

Prefix	Meaning
mono-	1
di-	2
tri-	3
tetra-	4
penta-	5
hexa-	6
hepta-	7
octa-	8
nona-	9
deca-	10


32



## Molecular Compounds

HI	hydrogen iodide
NF <sub>3</sub>	nitrogen trifluoride
SO <sub>2</sub>	sulfur dioxide
N <sub>2</sub> Cl <sub>4</sub>	dinitrogen tetrachloride
NO <sub>2</sub>	nitrogen dioxide
N <sub>2</sub> O	dinitrogen monoxide


33




An **acid** can be defined as a substance that yields hydrogen ions ( $\text{H}^+$ ) when dissolved in water.

For example: HCl gas and HCl in water

•Pure substance, hydrogen chloride HCl



•Dissolved in water ( $\text{H}_3\text{O}^+$  and  $\text{Cl}^-$ ), hydrochloric acid



35

**TABLE 2.5** Some Simple Acids


Anion	Corresponding Acid
$\text{F}^-$ (fluoride)	HF (hydrofluoric acid)
$\text{Cl}^-$ (chloride)	HCl (hydrochloric acid)
$\text{Br}^-$ (bromide)	HBr (hydrobromic acid)
$\text{I}^-$ (iodide)	HI (hydroiodic acid)
$\text{CN}^-$ (cyanide)	HCN (hydrocyanic acid)
$\text{S}^{2-}$ (sulfide)	$\text{H}_2\text{S}$ (hydrosulfuric acid)

36



An **oxoacid** is an acid that contains hydrogen, oxygen, and another element.



nitric acid



carbonic acid




phosphoric acid



37

### Naming Oxoacids and Oxoanions



38

The rules for naming **oxoanions**, *anions of oxoacids*, are as follows:

1. When all the H ions are removed from the “-ic” acid, the anion’s name ends with “-ate.”
2. When all the H ions are removed from the “-ous” acid, the anion’s name ends with “-ite.”
3. The names of anions in which one or more but not all the hydrogen ions have been removed must indicate the number of H ions present.

For example:

- $\text{H}_2\text{PO}_4^-$  dihydrogen phosphate
- $\text{HPO}_4^{2-}$  hydrogen phosphate

39

**TABLE 2.6** Names of Oxoacids and Oxoanions That Contain Chlorine

Acid	Anion
$\text{HClO}_4$ (perchloric acid)	$\text{ClO}_4^-$ (perchlorate)
$\text{HClO}_3$ (chloric acid)	$\text{ClO}_3^-$ (chlorate)
$\text{HClO}_2$ (chlorous acid)	$\text{ClO}_2^-$ (chlorite)
$\text{HClO}$ (hypochlorous acid)	$\text{ClO}^-$ (hypochlorite)

40

A **base** can be defined as a substance that yields hydroxide ions ( $\text{OH}^-$ ) when dissolved in water.

$\text{NaOH}$  sodium hydroxide

$\text{KOH}$  potassium hydroxide

$\text{Ba}(\text{OH})_2$  barium hydroxide

41


**Hydrates** are compounds that have a specific number of water molecules attached to them.

$\text{BaCl}_2 \cdot 2\text{H}_2\text{O}$  barium chloride dihydrate

$\text{LiCl} \cdot \text{H}_2\text{O}$  lithium chloride monohydrate

$\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$  magnesium sulfate heptahydrate


$\text{Sr}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$  strontium nitrate tetrahydrate




42

**Organic chemistry** is the branch of chemistry that deals with carbon compounds


### Functional Groups




methanol




CH<sub>3</sub>OH




methylamine



CH<sub>3</sub>NH<sub>2</sub>



acetic acid



CH<sub>3</sub>COOH

43