

Chem 1141

Fall 2012

Exam 3A

Name: _____

Please write your full name, and which exam version (3A) you have on the scantron sheet.

Multiple Choice. [3 points each.] Record your answers to the multiple choice questions on the scantron sheet.

Q1. Which element exists as a diatomic gas in its most stable form at 1 atm and 25 °C?

- a) helium b) carbon c) sulfur d) nitrogen e) argon

Q2. The pressure of a gas at STP is:

- a) 1 atm b) 1 Pa c) 1 mmHg d) 1 torr e) 1 N

Q3. The law that states that the pressure of a gas is inversely proportional to volume is:

- a) Avogadro's b) Boyle's c) Charles' d) Gay Lussac's e) Newton's


Q4. In the van der Waals equation for a gas, what does the constant b account for?

- a) The tendency of the molecules to stick together
b) The molecules can be cooled down to form a liquid
c) The temperature conversion from degrees celcius to Kelvin
d) The molecules are not all diatomic
e) The molecules have size

Q5. Given the following thermochemical equation:



calculate the value of ΔH° for the following reaction:




- a) -6.01 kJ/mol b) -12.02 kJ/mol c) +6.01 kJ/mol
d) +12.02 kJ/mol e) 36.1 kJ/mol

Q6. A reaction with a negative value of ΔH is said to be:

- a) Exogonic b) Endergonic c) Exothermic d) Endothermic

Q7. Given the thermochemical equation:



Then how much heat will be released when 3 mol of C_2H_2 is burned?

- a) 2598.8 kJ b) 866 kJ c) 7796.4 kJ d) 1732.5 kJ e) 3898.2 kJ

Q8. The reaction corresponding to the standard enthalpy of formation of trinitrotoluene, $C_6H_3N_3O_3(s)$ is:

- a) $C_6H_3N_3O_3(s) \rightarrow 6C(s) + 3H_2(g) + 3N_2(g) + 3O_2(g)$
- b) $6C(s, \text{graphite}) + H_2(g) + N_2(g) + O_2(g) \rightarrow C_6H_3N_3O_3(s)$
- c) $6C(s, \text{graphite}) + \frac{3}{2} H_2(g) + \frac{3}{2} N_2(g) + O_2(g) \rightarrow C_6H_3N_3O_3(s)$
- d) $6C(s, \text{graphite}) + \frac{3}{2} H_2(g) + \frac{3}{2} N_2(g) + \frac{3}{2} O_2(g) \rightarrow C_6H_3N_3O_3(s)$
- e) $12C(s, \text{graphite}) + 3 H_2(g) + 3 N_2(g) + 3 O_2(g) \rightarrow 2 C_6H_3N_3O_3(s)$

Q9. The term given to a particle of light is a(n):

- a) Wave
- b) Particle
- c) proton
- d) electron
- e) photon

Q10. The expression used to calculate the probability of an electron in space:

- a) ψ
- b) ψ^2
- c) ψ^3
- d) ψ^4
- e) $1/\psi$

Q11. The name given to the quantum number, l

- a) principal quantum number
- b) electron-spin quantum number
- c) angular momentum quantum number
- d) magnetic quantum number

Q12. The electron configuration for an atom of Cr is:

- a) $[He] 2s^2 2p^2$
- b) $[Ne] 3d^4$
- c) $[Ar] 4s^2 3d^4$
- d) $[Ar] 4s^1 3d^5$
- e) $[Ar] 3d^7$

Q13. The principle that says that the electrons in an atom prefer to enter orbitals in the same subshell with parallel spins, before pairing up:

- a) Hund's rule
- b) Paramagnetic rule
- c) Pauli's exclusion rule
- d) Bohr's condition
- e) Schrödinger's equation

Q14. What is the oxidation number of N in the compound: KNO_2 ?

- a) +5
- b) +3
- c) -1
- d) -3
- e) -5

Q15. Which SI prefix means $\times 10^{-9}$?

- a) μ
- b) m
- c) k
- d) n
- e) f

Short Response.

Show all work to receive credit. You must use the factor-label (conversion-factor) method for all conversions. Be sure to show all units and write your answers using the correct number of significant figures or decimal places.

Q16. [8 pts.] Write the full electron configuration and orbital diagrams for the following atoms:


a) Al

b) Cu

Q17. [8 pts.] Given the following data:

$\Delta H_f^\circ / \text{kJ} \cdot \text{mol}^{-1}$	+226.6	0	-393.5	-241.8
Compound	C ₂ H ₂ (g)	O ₂ (g)	CO ₂ (g)	H ₂ O(g)

(i) Predict ΔH° for the following reaction:



(ii) Calculate how much heat is absorbed/released (state which!) if 12.0 g of C₂H₂(g) and 15.0 g of O₂(g) react.

Q18. [8 pts.] Give a detailed explanation of how *real* gases differ from *ideal* gases.

Q19. [8 pts.] 5.0 g of Ar(g) and 5.0 g of Ne(g) is released into an empty 5.0-L container at a temperature of 17 °C. Calculate the partial pressures of each gas, the mole-fractions of each gas, and the total pressure inside the container.

Q20. [8 pts.] A 4.51 g sample of a metal at 182 °C is dropped into a calorimeter containing 43.2 g of water at 5.1 °C. Given that the water has a specific heat of 4.184 J/g·°C, and assuming that the calorimeter forms a perfectly isolated system, calculate the specific heat of the metal if the final temperature of the system is 8.4 °C.

Q21. [5 pts.] Calculate the empirical formula of a substance containing 40.1 percent carbon, 6.6 percent hydrogen, and 53.3 percent oxygen by mass.

Q22. [10 pts.] Write the molecular, full-ionic, and net-ionic equation for the reaction between aqueous sodium carbonate and aqueous magnesium nitrate.


Molecular:

Full-Ionic:

Net-Ionic:

BONUS Question

Sketch and label the five different 3d orbitals.



Quantum Mechanics

Partial List of Solubility Rules

TABLE 4.2 Solubility Rules for Common Ionic Compounds in Water at 25°C

Soluble Compounds	Exceptions
Halides (Cl ⁻ , Br ⁻ , I ⁻)	Halides of Ag ⁺ , Hg ₂ ²⁺ , and Pb ²⁺
Sulfates (SO ₄ ²⁻)	Sulfates of Ag ⁺ , Ca ²⁺ , Sr ²⁺ , Ba ²⁺ , Hg ₂ ²⁺ , and Pb ²⁺
Insoluble Compounds	Exceptions
Carbonates (CO ₃ ²⁻), phosphates (PO ₄ ³⁻), chromates (CrO ₄ ²⁻), and sulfides (S ²⁻)	Compounds containing alkali metal ions and the ammonium ion
Hydroxides (OH ⁻)	Compounds containing alkali metal ions and the Ba ²⁺ ion

Useful Information:

$$pV = nRT \quad \left(p + \frac{an}{V^2}\right)(V - nb) = nRT \quad 1 \text{ atm} = 760 \text{ mmHg} = 101325 \text{ Pa} \quad R = 0.08206 \frac{\text{atm} \cdot \text{L}}{\text{mol} \cdot \text{K}}$$

$$M_1V_1 = M_2V_2 \quad N_A = 6.022 \times 10^{23} \quad q = m \cdot s \cdot \Delta t \quad q = C \cdot \Delta t$$

$$c = v\lambda \quad E = h\nu \quad c = 3.00 \times 10^8 \text{ m/s} \quad h = 6.626 \times 10^{-34} \text{ J} \cdot \text{s}$$

$$E_n = -R_H \left(\frac{1}{n^2}\right) \quad R_H = 2.18 \times 10^{-18} \text{ J} \quad \lambda = \frac{h}{mu}$$

Periodic Table

1 IA	2 IIA											13 IIIA	14 IVA	15 VA	16 VIA	17 VIIA	18 VIIIA	
1 H 1.01																		2 He 4.00
3 Li 6.94	4 Be 9.01											5 B 10.81	6 C 12.01	7 N 14.01	8 O 16.00	9 F 19.00	10 Ne 20.18	
11 Na 22.99	12 Mg 24.31	3 IIIB	4 IVB	5 VB	6 VIB	7 VIIB	8	9 VIIIB	10	11 IB	12 IIB	13 Al 26.98	14 Si 28.09	15 P 30.97	16 S 32.07	17 Cl 35.45	18 Ar 39.95	
19 K 39.1	20 Ca 40.08	21 Sc 44.96	22 Ti 47.88	23 V 50.94	24 Cr 52.00	25 Mn 54.94	26 Fe 55.85	27 Co 58.93	28 Ni 58.69	29 Cu 63.55	30 Zn 65.39	31 Ga 69.72	32 Ge 72.61	33 As 74.92	34 Se 78.96	35 Br 79.90	36 Kr 83.80	
37 Rb 85.47	38 Sr 87.62	39 Y 88.91	40 Zr 91.22	41 Nb 92.91	42 Mo 95.94	43 Tc (98)	44 Ru 101.07	45 Rh 102.91	46 Pd 106.42	47 Ag 107.87	48 Cd 112.41	49 In 114.82	50 Sn 118.71	51 Sb 121.76	52 Te 127.6	53 I 126.9	54 Xe 131.29	
55 Cs 132.9	56 Ba 137.3	57 La* 138.9	72 Hf 178.5	73 Ta 180.9	74 W 183.9	75 Re 186.2	76 Os 190.2	77 Ir 192.2	78 Pt 195.1	79 Au 197.0	80 Hg 200.6	81 Tl 204.4	82 Pb 207.2	83 Bi 209	84 Po (209)	85 At (210)	86 Rn (222)	
87 Fr (223)	88 Ra (226)	89 Ac^ (227)	104 Rf (261)	105 Db (262)	106 Sg (263)	107 Bh (264)	108 Hs (265)	109 Mt (268)	110 Ds (271)	111 Rg (272)								

* 58 Ce 140.1	59 Pr 140.9	60 Nd 144.2	61 Pm (145)	62 Sm 150.4	63 Eu 152.0	64 Gd 157.3	65 Tb 158.9	66 Dy 162.5	67 Ho 164.9	68 Er 167.3	69 Tm 168.9	70 Yb 173.0	71 Lu 175.0
^ 90 Th 232.0	91 Pa (231)	92 U 238.0	93 Np (237)	94 Pu (244)	95 Am (243)	96 Cm (247)	97 Bk (247)	98 Cf (251)	99 Es (252)	100 Fm (257)	101 Md (258)	102 No (259)	103 Lr (260)