# **General Chemistry 2 (CHEM 1141)**

# Shawnee State University - Autumn 2022

October 20, 2022

# Exam # 2A

Name \_\_\_\_

•	Please print your full name, and the exam version (2A) that you have on the scantron sheet!  (Bubble in the best answer choice for each question on the scantron sheet in pencil!)										
Please ☑ check t	the box next to your correct section num	ber.									
Section #:	☐ 1. (Mon Lab, 10:10 AM – 1:00 PM) ☐ 2. (Wed Lab, 10:10 AM – 1:00 PM) ☐ 3. (Tue Lab, 11:00 AM – 1:50 PM) ☐ 4. (Thu Lab, 11:00 AM – 1:50 PM)	Fleeman   Napper									

Multiple Choice: \_\_\_\_\_ **/ 50** Q21: \_\_\_\_\_ / 10 Q22: \_\_\_\_\_ / 10 Q23: \_\_\_\_\_ / 10 Q24: \_\_\_\_\_ / 10 Q25: \_\_\_\_\_ / 10 **BONUS:** \_\_\_\_\_ /3 **TOTAL:** \_\_\_\_\_ / 100



### Each problem in this section (multiple choice) is worth 2.5 points!



Q1. The volume of a gas is proportional to the temperature of a gas is known as

- A) Avogadro's Law
- B) Charles's Law
- C) Boyle's Law
- D) Ideal Gas Law

Q2. Identify the major ionic species present in an aqueous solution of FeCl<sub>3</sub>:

- A) Fe+, 3 Cl-
- B) Fe<sup>2+</sup>, 3 Cl<sup>-</sup>
- C) Fe<sub>3+</sub>, 3 Cl-
- D) Fe<sup>3+</sup>, 3 Cl<sup>3-</sup>

Q3. For which of the following acids is chlorine in the +5 oxidation state?

- A) HClO<sub>2</sub>
- B) HClO<sub>3</sub>
- C) HClO<sub>4</sub>
- D) HCl

Q4. How many moles of water are produced when 1.0 mole of NH<sub>3</sub> reacts according to the following chemical equation?

$$4~NH_3 + 5~O_2 \longrightarrow ~4~NO + 6~H_2O$$

- A) 0.67 moles
- B) 1.0 moles
- C) 1.3 moles
- D) 1.5 moles

- Q5. A 20.00 mL sample of 0.1015 M nitric acid is introduced into a flask, and water is added until the volume of the solution reaches 250. mL. What is the concentration of nitric acid in the final solution?
  - A)  $3.25 \times 10^{-2}$  M
  - B)  $8.12 \times 10^{-3} \text{ M}$
  - C) 0.406 M
  - D)  $5.08 \times 10^{-4}$
- Q6. A mixture of three gases has a total pressure of 1.82 atm at 298 K. The mixture is found to contain 1.27 mol CO<sub>2</sub>, 3.04 mol CO, and 1.50 mol Ar.

What is the partial pressure of Ar?

- A) 0.258 atm
- B) 1.50 atm
- C) 0.470 atm
- D) 0.824 atm
- Q7. A volume of gas occupies  $1.40 \times 10^3$  mL at 25 °C and 760 mmHg. What volume will it occupy at the same temperature and 380 mmHg?
  - A) 700 mL
  - B) 1,050 mL
  - C) 1,140 mL
  - D) 2,800 mL
- Q8. Choose the statement below that is **false**:
  - A) a weak acid solution consists mostly of non-ionized acid molecules
  - B) non-electrolyte solutions do not conduct electricity
  - C) a strong acid solution consists of only partially ionized acid molecules
  - D) a strong electrolyte completely dissociates into ions in water

Q9. The following equation represents the formation of nitrogen dioxide, a major component of smog:

$$2 \text{ NO} + \text{O}_2 \longrightarrow 2 \text{ NO}_2$$

If 0.88 mol of NO is reacted with 0.79 mol of  $O_2$  to produce  $NO_2$ , the limiting reactant is

- A) NO
- B)  $O_2$
- C) NO<sub>2</sub>
- D) both NO and O2
- Q10. You have two HCl solutions, labeled solution A and solution B. Solution A has a greater concentration than solution B. Which of the following statements is true?
  - A) If you have equal volumes of both solutions, solution B must contain more moles of HCl
  - B) If you have equal moles of HCl in both solutions, solution B must have a greater volume.
  - C) To obtain equal concentrations of both solutions, you must add a certain amount of water to solution B.
  - D) Adding more moles of HCl to both solutions will make them less concentrated.
- Q11. Which combination of solutions will result in a solid being formed?
  - A)  $AgNO_3(aq) + HCl(aq)$
  - B)  $HBr(aq) + LiHCO_3(aq)$
  - C)  $NaCl(aq) + Na_2CO_3$
  - D)  $NH_4NO_3(aq) + KI(aq)$
- Q12. Which substance would cause litmus to turn blue?
  - A) NaCl(aq)
  - B) NH<sub>4</sub>NO<sub>3</sub>(aq)
  - C) HC<sub>2</sub>H<sub>3</sub>O<sub>2</sub>(aq)
  - D) LiOH(aq)

- Q13. 13.8 g of LiBr(s) is dissolved in water, so that the total volume is 250. mL. What is the molar concentration of the solute?
  - A) 0.159 M
  - B) 0.636 M
  - C) 0.0006 M
  - D) 0.0552 M
- Q14. For the following redox reaction:

$$2Fe^{2+}(aq) + 2H^{+}(aq) + H_2O_2(aq) \rightarrow 2Fe^{3+}(aq) + 2H_2O(l)$$

The <u>reducing</u> agent is

- A) Fe<sup>2+</sup>
- B)  $H_2O_2$
- C) H<sup>+</sup>
- D) Fe<sub>3+</sub>
- Q15. Calculate the density of  $CO_2(g)$  in the atmosphere of Venus, where the temperature is 467 °C at a pressure of 93 atm.
  - A) 67.4 g/L
  - B)  $3.69 \times 10^7 \text{ g/L}$
  - C) 107 g/L
  - D)  $2.33 \times 10^7 \, g/L$
- Q16. How many moles of LiBr are contained in 20.00 mL of 0.500 M LiBr(aq)?
  - A) 10.0 mol
  - B) 0.010 0 mol
  - C) 40.0 mol
  - D) 0.025 0 mol

- Q17. When dealing with gases, standard temperature and pressure corresponds to:
  - A) 1 atm, o °C
  - B) 1 atm, 100 °C
  - C) 760 mmHg, 0 K
  - D) 760 mm Hg, 100 K
- Q18. Which of the following elements is a diatomic gas at room temperature and pressure?
  - A) argon
  - B) boron
  - C) carbon
  - D) chlorine
- Q19. Which net ionic equation cannot be correct?
  - A)  $Ag^{+}(aq) + Cl^{-}(aq) \rightarrow AgCl(s)$
  - B)  $H^+(aq) + OH^-(aq) \rightarrow H_2O(l)$
  - C)  $Na_2^+(aq) + Cl_2^-(aq) \rightarrow Na_2Cl_2(s)$
  - D)  $3Ca^{2+}(aq) + 2PO_4^{3-}(aq) \rightarrow Ca_3(PO_4)_2(s)$
- Q20. What is the best piece of glassware to use when preparing a solution of known concentration in the lab?
  - A) Erlenmeyer flask
  - B) graduated cylinder
  - C) beaker
  - D) volumetric flask



Each problem in this section (short answer) is worth 10 points!

All work must be shown to receive credit!

You must use the factor–label (conversion–factor) method for all conversions!

Be sure to include units where applicable!

All numeric answers must be rounded to the correct number of significant figures!




Q21. Provide a correctly balanced reaction equation that shows the chemical reaction that takes place when an aqueous solution of iron(II) bromide is mixed with an aqueous solution of potassium carbonate. In addition, provide the correct full ionic as well as the net ionic equation for this reaction. **Be sure to show all state symbols and charges where appropriate.** 

Complete Balanced	Reaction	<b>Equation</b>	(Molecular	<b>Equation</b> )

$\_$ FeBr <sub>2</sub> (aq) + $\_$ K <sub>2</sub> CO <sub>3</sub> (aq) $\rightarrow$	+	

**Complete Ionic Equation** 

**Net Ionic Equation** 



Q23. A 25.00 mL sample of battery acid (H<sub>2</sub>SO<sub>4</sub>) is obtained and placed in a flask, along with two drops of phenolphthalein indicator. A buret is filled with 0.100 0 M NaOH(aq) and placed above the acid. If the initial reading of the buret is 0.03 mL, and the final reading (when a pale pink color persists) is 32.13 mL—calculate the molar concentration (molarity) of the acid.

Show all work, being sure to include units and the correct number of digits at every step in your calculation.

$$H_2SO_4(aq) + 2 NaOH(aq) \rightarrow 2 H_2O(l) + Na_2SO_4(aq)$$

Q24.	(a) 100.0 mL of 3.00 M HCl(aq) was added to an excess of Na <sub>2</sub> CO <sub>3</sub> (aq). What volume
	of gas should be formed at a temperature of 23 °C and a pressure of 753 mmHg?

$$2 \text{ HCl(aq)} + \text{Na}_2\text{CO}_3(\text{aq}) \rightarrow 2 \text{ NaCl(aq)} + \text{H}_2\text{O(l)} + \text{CO}_2(\text{g})$$

(b) If the actual volume of gas formed was 3.37 L—calculate the percent yield of the reaction.

Q25.	The production of aluminum from bauxite is an incredibly energy intensive process. Approximately 5% of electricity generated in the United States is used to produce it.
	The simplified chemical equation for this process is:

$$2 \text{ Al}_2\text{O}_3(1) \rightarrow 4 \text{ Al}(1) + 3 \text{ O}_2(g)$$

Being sure to use the conversion-factor method, showing all work, and using the correct number of digits, please answer the following questions:

(a) If 0.100 mol of oxygen is formed, how many moles of aluminum oxide must have been used up?

(b) If 2,240 g of Al<sub>2</sub>O<sub>3</sub> are used up, how many moles of aluminum can be made?

(c) If 2,240 g of  $Al_2O_3$  are used up, and 1090 g of aluminum are formed, calculate the percent yield of this reaction.

# 

Name three strong inorganic acids	Name t	ne thre	ee stro	ng inor	ganic	acids.
-----------------------------------	--------	---------	---------	---------	-------	--------

1.				
2.				

3.\_\_\_\_\_

# **Exam checklist:**

#### (Check the boxes to certify the following:)

- ☐ My correct lab section has been indicated on the front page
- ☐ My full name is written legibly on the scantron sheet
- ☐ My exam version (A, B, C, or D) is written on the scantron sheet
- ☐ I have shown work for all problems (where appropriate), paying attention to
  - o Significant figures / decimal places
  - o Units
- ☐ I have used the conversion-factor method for all conversions
- ☐ If I have torn off the back page (periodic table), I will not turn it in with my exam!

Thank you from the Chemistry Professors and Good Luck!



# **TABLE 4.2** Solubility Rules for Common Ionic Compounds in Water at 25°C

Soluble Compounds	Exceptions
Halides (Cl <sup>-</sup> , Br <sup>-</sup> , I <sup>-</sup> ) Sulfates (SO <sub>4</sub> <sup>2-</sup> )	Halides of $Ag^+$ , $Hg_2^{2+}$ , and $Pb^{2+}$ Sulfates of $Ag^+$ , $Ca^{2+}$ , $Sr^{2+}$ , $Ba^{2+}$ , $Hg_2^{2+}$ , and $Pb^{2+}$
Insoluble Compounds	Exceptions
Carbonates $(CO_3^{2-})$ , phosphates $(PO_4^{3-})$ , chromates $(CrO_4^{2-})$ , and sulfides $(S^{2-})$	Compounds containing alkali metal ions and the ammonium ion
Hydroxides (OH <sup>-</sup> )	Compounds containing alkali metal ions and the Ba <sup>2+</sup> ion

# **Useful information:**

$$M_1V_1 = M_2V_2$$

$$N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$$

1 atm = 101,325 Pa = 760 mmHg = 760 torr

 $R = 0.08206 \text{ L} \cdot \text{atm/mol} \cdot \text{K}$ 

$$p_1V_1 = p_2V_2$$

$$\frac{V_1}{T_1} = \frac{V_2}{T_2}$$

$$\frac{p_1 V_1}{T_1} = \frac{p_2 V_2}{T_2}$$

$$pV = nRT$$

$$p\mathcal{M} = dRT$$

VIIIA 81	<b>F</b> 5	4.003	10	Se	20.18	18	Ā	39.95	36	궃	83.80	54	Xe	131.3	98	R	[222]	118		[293]						
AII V		17	6	ш	19.00	17	ਠ	35.45	35	Б	79.90	53	_	126.9	85	¥	[210]	117								
VIA		16	8	0	16.00	16	ഗ	32.07	34	Se	78.96	52	Te	127.60	84	Ъ	[210]	116		[289]	70	Ϋ́	173.0	102	å	[259]
<b>X</b>		15	2	z	14.01	15	۵	30.97	33	As	74.92160	51	Sb	121.8	83	Bi	209.0	115			69	Tm	168.9	101	Md	[258]
ĕ		14	9	ပ	12.01	14	S	28.09	32	Ge	72.61	20	Sn	118.7	82	Pb	207.2	114		[285]	89	Щ	167.3	100	Fm	[257]
<b>ĕ</b>		13	9	В	10.81	13	₹	26.98	31	Ga	69.72	49	므	114.8	81	F	204.4	113			29	P H	164.9	66	Es	[252]
								12	30	Zn	62.39	48	ပ်	112.4	80	H	200.6	112		[277]	99	ò	162.50	86	ర	[251]
								11	59	Cn	63.55	47	Ag	107.9	79	Αn	197.0	111		[272]	65	q L	158.9	26	ਲ	[247]
ents								10	28	Z	58.69	46	Pd	106.4	78	Ŧ	195.1	110		[269]	64	Вq	157.3	96	S	[247]
Elem								6	27	ပိ	58.93	45	格	102.9	77	<u>_</u>	192.2	109	Ĭ	[268]	63	Eu	152.0	98	Am	[243]
of the								89	56	Бe	55.85	44	Ru	101.1	9/	Os	190.2	108	Hs	[265]	62	Sm	150.4	94	Pu	[244]
Periodic Table of the Elements								7	25	Σ	54.94	43	ည	[86]	75	Re	186.2	107	Bh	[264]	61	Pm	[145]	93	S N	[237]
dic T								9	24	ပ်	52.00	42	Θ	95.94	74	>	183.8	106	Sg	[266]	09	Š	144.2	92	_	238.0
Perio								2	23	>	50.94	41	Q N	92.91	73	Та	180.9	105	Ob	[262]	29	Ā	140.9	91	Ра	231.0
								4	22	F	47.87	40	Zr	91.22	72	¥	178.5	104	Ŗ	[261]	28	င္ပ	140.1	06	ᄕ	232.0
								က	21	လွ	44.96	39	>	88.91	7.1	Ľ	175.0	103	ב	[262]	22	La	138.9	88	Ac	[227]
¥		2	4	Be	9.012	12	Mg	24.31	20	Ca	40.08	38	ร	87.62	56	Ba*	137.3	88	Ra∗∗	[226]		*			*	
₫ -	- <b>I</b>	1.008	8	=	6.941	11	Na	22.99	19	¥	39.10	37	8 e	85.47	55	S	132.9	87	ቷ	[223]						